

AD-A173 899

THEORETICAL STUDIES RELATING TO THE INTERACTION OF
RADIATION WITH MATTER... (U) NEW YORK UNIV NY DEPT OF
PHYSICS P R BERMAN 01 OCT 86 N00014-77-C-0553


1/1

UNCLASSIFIED

F/G 11/6

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A173 899

12

Annual Report (AR9)

Title: "Theoretical Studies Relating to the Interaction of
Radiation with Matter: Atomic Collision Processes
Occurring in the Presence of Radiation Fields"

Supported by: The U.S. Office of Naval Research

Contract No: N00014-77-C-0553

Report Period: August 1, 1985 - July 31, 1986

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

Approved for public release; distribution unlimited.

DTIC
ELECTE
OCT 29 1986
S B

DTIC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AR 9	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <i>THEORETICAL STUDIES RELATING TO THE</i> Interaction of Radiation with Matter: Atomic Collision Processes Occurring in the Presence of Radiation Fields"		5. TYPE OF REPORT & PERIOD COVERED Interim 8/1/85 - 7/31/86
7. AUTHOR(s) P.R. Berman		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Prof. P.R. Berman Physics Dept. - New York University 4 Washington Place, New York, N.Y. 10003		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0553
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Resident Representative New York 715 Broadway, New York, N.Y. 10003		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61153N RR011-07-02 NR 396-029
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE October 1, 1986
		13. NUMBER OF PAGES 150
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Laser Spectroscopy, Optical Collisions, Radiative Collisions, Two-level Atom, Atomic Coherence, Degenerate Four-Wave Mixing, Saturation Spectroscopy, Laser Propagation, Dressed Atom, Photon Echo, Bloch Equations, Collision Kernel, Collisions, Optical Noise		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Work is reported in areas of:- (1) Creation of Electronic State Coherences via Laser-Assisted Collisions (2) Collision Effects in Four-Wave Mixing and Pump-Probe Spectroscopy (3) Modified Optical Bloch Equations in Solids (4) Transient Spectra in Atom-Field Interactions (5) Collision Kernels and Transport Coefficients CONTINUED ON BACK		

- (6) Collisions in Strong cw and Transient Laser Fields
- (7) Quantum vs. Classical Description of Laser Fields
- (8) Intense Optical Noise Sources
- (9) Laser-Assisted Collisions

Accession	
NTIS	✓
DTIC	
Uncl.	
Just.	
PER CALL JC	
By	
Dist	
Avail.	
Dist	
A-1	



Research has been carried out in the areas of (1) Modified Optical Bloch Equations in vapors and solids, (2) Creation of electronic-state coherence in laser-assisted collisions (3) Collisionally modified line shapes in pump-probe and 4-wave mixing spectroscopy, (4) Collision kernels and transport coefficients, (5) Collisions in strong cw and transient laser fields, (6) Quantum vs classical description of laser fields in nonlinear spectroscopy, (7) Intense optical noise sources, (8) Laser-assisted collisions and (9) Transient spectra in atom-field interactions.

1. Modified Optical Bloch Equations (P. Berman)


The work outlined in last year's annual report¹ has been completed. A "strong redistribution" model of Markovian fluctuations was used to analyze the Free-Induction Decay (FID) signal emitted by atoms which were subjected to random frequency fluctuations.² The results of this calculation helped to explain in a qualitative manner the experimental results of DeVoe and Brewer³ which had stimulated a great deal of theoretical activity in this area. A detailed examination of the validity requirements for the Optical Bloch Equations was presented in a second article.^{4*} In a companion article,^{5*} a general method was developed for calculating the effects of atomic frequency fluctuations in laser-atom interactions. This method is particularly useful if the atom-field interaction can be treated in a perturbative expansion. Explicit time-dependent and time-independent frequency fluctuation propagators were derived for (a) strong redistribution, (b) "difference kernel" and (c) Brownian motion models for the frequency fluctuations. It was shown that no consistent explanation of the DeVoe-Brewer results seems to exist. The next step in this problem would be to extend the method to include saturation effects; this extension represents an extremely difficult task.

2. Creation of Electronic-State Coherence in Collisionally-Assisted Collisions (P. Berman)

Experimental evidence has been obtained for the creation of collision-

* An asterisk indicates that a reprint or preprint of this article has been forwarded to the Scientific Officer with this report. Reprints of articles have been furnished to DTIC with this report. Preprints or reprints of these articles are available on request to anyone receiving this report.

induced, optical coherence in Na.^{6*} The reaction that was studied is



where the notation $(4D-3P_{3/2})$ indicates a coherent superposition of the 4D and $3P_{3/2}$ states. The laser at frequency Ω_1 is tuned about 200 GHz below the $3S-3P_{3/2}$ transition frequency. When the second laser's frequency Ω_2 is scanned, a resonance structure is observed when $2\Omega_2$ is detuned from the $3S-4D$ transition by the same amounts as the first laser. This resonance structure is observed in emission at the $4D-3P_{3/2}$ transition frequency (essentially via a 4-wave mixing process), but is absent in the absence of collisions. Thus, it has been demonstrated that it is possible to generate coherent optical radiation in a collision-induced reaction.

3. Collisionally-Modified Line Shapes in Pump-Probe and 4-wave Mixing Spectroscopy (G. Khitrova, P.R. Berman)

G. Khitrova has finished her thesis in which new features of the line shapes in pump-probe spectroscopy and 4-wave mixing on a single atomic transition are examined. In particular, new effects related to (a) effects of magnetic degeneracy, (b) velocity-changing collisions and (c) contributions from all velocity subclasses of atoms (and not from only those Doppler-shifted into resonance with the fields) have been reported. The conditions under which narrow resonances (characterized by the ground state width) can be observed are clearly established. A short summary of the various topics is contained in last year's Annual Report.¹

4. Collision Kernels and Transport Coefficients (P. Berman)

In collaboration with the group of J. Woerdman (University of Leiden), we have established an integral relation between the collision kernels used in analyzing laser spectroscopy experiments and the transport coefficients of classical transport theory.^{8*} Our results are extremely useful in the analysis of laser spectroscopy experiments involving collisions. By using experimental or theoretical values for the various

transport coefficients, one can reduce the number of free parameters needed to fit laser spectroscopic line shapes in which collisions play a role. This method has been applied to a study of Na - rare gas collisions, observed using saturation spectroscopy. Good fits to the data were obtained with essentially no free parameter. A knowledge of collisional parameters for the Na - rare gas system is critical for understanding the theory of the so-called "optical piston".⁹

5. Collisions in Strong cw and Transient Laser Fields (P. Berman)

Attempts continue to obtain "propagators" which give the time-development for a "two-level" atom subject to both a strong external field and atomic phase and frequency fluctuations. The original motivation for this problem is linked to an experiment¹⁰ in Mossberg's group in which he claimed quenching of the effects of velocity-changing collisions at high laser field strengths in a photon echo experiment. As yet, there has been no rigorous proof of that conjecture. Our calculation is intended to supply an answer to this problem. Some progress has been made using a dressed-atom approach to the problem, but we are still unable to confirm or reject Mossberg's interpretation of his result. The problem we are studying is fundamental and the results, when obtained, can be applied to a wide variety of problems involving atoms in strong fields undergoing phase or frequency fluctuations.

6. Quantum vs Classical Descriptions of Laser Fields in Nonlinear Spectroscopy (P. Berman)

In collaboration with G. Grynberg, we have started to examine some aspects of nonlinear spectroscopy related to the quantized nature of the electromagnetic field. The motivation for this study is based on obtaining a physical understanding of "collision-induced extra resonances" in 4-wave mixing experiments.¹¹ In such experiments, it is often stated that collisions lead to a "real" (i.e. non-virtual) excitation of level coherence. If the level coherence in such a problem is calculated using a fixed number quantized radiation field, it can be shown that the level coherence vanishes identically. Does this mean that the pressure-induced resonance

resonance would vanish if fixed number states of the radiation field were used experimentally? To answer this question, we have begun a systematic study of various experimental situations of current interest to determine which of these experiments would provide the most sensitive probe of the quantum vs classical nature of the radiation field.

7. Intense Optical Noise Sources

Several experimental groups¹² are now using large-band laser sources as a means for obtaining femtosecond time resolution. The major idea behind the work is that one can exploit the short coherence time ($\tau_c = 1.0$ ps) of a large-band source to arrive at a time resolution which is of order τ_c . Stimulated photon echo experiments, have been carried out which have dramatically demonstrated the feasibility of this idea.¹² However, there have been field saturation effects which cannot be explained within the context of current theories. We have started to formulate a theory for the stimulated photon echo with broad-band noise sources. Some progress has been made analytically, but a large scale numerical solution may be necessary. The theoretical problem is complicated by the fact that the same noise is sent into an atomic vapor with a time delay. This means that the system has a memory and this memory greatly complicates the analysis.

8. Laser-assisted Collisions (P. Berman)

In collaboration with A. Bambini (Quantum Electronics Institute, Florence), we have tried to explain a long-standing mystery regarding the far wing behavior of LICET (laser-induced collisional energy transfer) profiles. Theory, based on an effective two-level problem, predicts a $\Delta^{-0.5}$ (Δ = detuning from resonance) dependence while experiment varies from $\Delta^{-0.5}$ to $\Delta^{-0.85}$. Improving the existing calculations by including some additional contributions from a third (virtual) level has enabled us to give a unified explanation of existing experiments. An article is in preparation.

In collaboration with F. Schuller (High-Pressure Laboratory - Villetaneuse, France) we are trying to extend a previous calculation¹³ on

effects of magnetic degeneracy in LICET into the line wing.

9. Transient Spectra in Atom-Field Interactions

In collaboration with the group of T. Mossberg (Harvard), we have studied the transient spectra produced when atoms are prepared in pure dressed states of the atom and radiation field. Both transient Autler-Townes spectra^{14*} and transient resonance fluorescence^{15*} for atoms prepared in dressed states have been studied theoretically. It was shown that one of the peaks in the Autler-Townes doublet and fluorescence triplet is suppressed at early times. A simple explanation, based on a dressed-atom picture, was used to understand the results. Experimental verification of the transient Autler-Townes doublet for pure dressed-state preparation has been achieved.^{16*} Theoretical calculations for the transient pump-probe spectroscopy of a single transition is in progress.

10. Miscellaneous

A previous review article on optical pumping in nonlinear spectroscopy has appeared as has our work on photon echoes using double-resonance pulses.^{17*,18*} Previous works on two-level systems^{19*,20*} corrections to exponential decay produced by adiabatic switching^{21*} and a rederivation of Wigner's infinite product representation of the R matrix^{22*} were published. The effect of an unstable continuum on the exponential decay law was studied.^{23*} A calculation relating to the treatment of intermediate states in constructing Green function operators was carried out.^{24*} An analogy between certain integrals appearing in the theory of resonance fluorescence following pulse excitation and certain diffraction integrals was obtained.^{25*} Finally, some earlier work on pulse propagation involving the interaction of two fields with a "three-level" atom has appeared.^{26*}

References

1. Annual Report AR 8
2. P.R. Berman and R.G. Brewer, Phys. Rev. A32, 2784 (1985).
3. R.G. Devoe and R.G. Brewer, Phys. Rev. Lett. 50, 1269 (1983).
4. P.R. Berman, J. Opt. Soc. Amer. B3, 564 (1986); see also P.R. Berman and R.G. Brewer in Laser Spectroscopy VII, edited by T.W. Hänsch and Y.R. Shen (Springer-Verlag, Berlin, 1985) pp. 270-273. This article also appears in Methods of Laser Spectroscopy, edited by Y. Prior, A. Ben Reuven and M. Rosenbluh (Plenum, New York, 1986) pp. 205-208.
5. P.R. Berman, J. Opt. Soc. Amer. B3, 572 (1986).
6. E. Giacobino and P.R. Berman, Phys. Rev. Lett. (submitted).
7. G. Khitrova, Thesis, 1986, unpublished.
8. P.R. Berman, J.E.M. Haverkort and J. Woerdman, Phys. Rev. A (to appear).
9. See, for example, H.G.C. Werij, J.E.M. Haverkort, and J.P. Woerdman, Phys. Rev. A33, 3270 (1986).
10. A.G. Yodh, J. Golub, N.W. Carlson and T.W. Mossberg, Phys. Rev. Lett. 53, 659 (1984).
11. See, for example, L.J. Rothberg and N. Bloembergen, in Laser Spectroscopy VI, edited by H.P. Weber and W. Luthy (Springer-Verlag, Berlin (1983) pp. 178-182 and references therein.
12. J.C. Keller and J.L. LeGouët, Phys. Rev. A32, 1624 (1985); S. Asaka, H. Nakatsuka, M. Fujiwara and M. Matsuoka, Phys. Rev. A29, 2286 (1984); D. DeBeer, L.G. Van Wagenen, R. Beach and S.R. Hartmann, Phys. Rev. Lett. 56, 1128 (1986); J.E. Golub and T.W. Mossberg, Opt. Lett. 11, 431 (1986).
13. P.R. Berman, Phys. Rev. A22, 1838, 1848 (1980).

References - Con't.

14. N. Lu, P.R. Berman, A.G. Yodh, Y.S. Bai and T.W. Mossberg, Phys. Rev. A33, 3956 (1986).
15. N. Lu, P.R. Berman, Y.S. Bai, J.E. Golub and T.W. Mossberg, Phys. Rev. A34, 319 (1986).
16. Y.S. Bai, T.W. Mossberg, N. Lu and P.R. Berman, Phys. Rev. Lett. 57, 1692 (1986).
17. P. R. Berman, Ann. Phys. (France) 10, 985 (1985).
18. N. Lu and P.R. Berman, J. Opt. Soc. Amer. B2, 1883 (1985).
19. E.J. Robinson, J. Phys. B18, 3687 (1985).
20. E.J. Robinson, J. Phys. B18, L657 (1985).
21. E.J. Robinson, Phys. Rev. A33, 1461 (1986).
22. E.J. Robinson, Phys. Rev. A34, 1896 (1986).
23. E.J. Robinson, Phys. Rev. Lett. 57, 1281 (1986).
24. E.J. Robinson, Phys. Lett., to appear.
25. E.J. Robinson, J. Phys. B, to appear.
26. F.P. Mattar, P.R. Berman, A.W. Matos, Y. Claude, C. Goutier and C.M. Bowden, in Multiple-Photon Excitation and Dissociation of Polyatomic Molecules, edited by C. Cantrell (Springer-Verlag, Berlin, 1986) pp. 223-283.

PUBLICATIONS

PAPERS SUBMITTED TO REFEREED JOURNALS
(Not yet published)

1. Y.S. Bai, T.W. Mossberg, N. Lu and P.R. Berman "Transient Suppression of the Autler-Townes Doublet", Phys. Rev. Letters (to appear) (additional support from NSF).
2. P.R. Berman, J.E.M. Haverkort and J.P. Woerdman, "Collision Kernels and Transport Coefficients," Physical Review A (to appear) (additional support from NSF).
3. E. Giacobino and P.R. Berman, "Collisionally-Aided Coherent Emission at an Optical Frequency," Phys. Rev. Letters (submitted) (additional support from NSF).
4. E.J. Robinson, "The Green's Function Method and the Pauli Principle in Intermediate States", Physics Letters A (1986), (to appear).
5. E.J. Robinson, "Temporal Diffraction and Eigenvalue Interpretation of the Resonance Fluorescence of Two-Level Systems Driven by Short Pulses". J. Phys. B (1986) (to appear).

PAPERS PUBLISHED IN REFEREED JOURNALS

1. P.R. Berman, "Current Research in Optical Pumping," Ann. Phys. (France) 10, 985-993 (1985).
2. N. Lu and P.R. Berman, "Photon Echoes Using Double-Resonance Pulses," J. Opt. Soc. Amer. B2, 1883-1893 (1985).
3. P.R. Berman and R.G. Brewer, "Modified Optical Bloch Equations for Solids", Phys. Rev. A32, 2784-2796 (1985).
4. P.R. Berman, "Validity Conditions for the Optical Bloch Equations," J. Opt. Soc. Amer. B3, 564-571 (1986).
5. P.R. Berman, "Markovian Relaxation Processes for Atoms in Vapors and Solids: Calculation of Free-Induction Decay in the Weak-External-Field Limit," J. Opt. Soc. Amer. B3, 572-586 (1986).
6. N. Lu, P.R. Berman, A.G. Yodh, Y.S. Bai, and T.W. Mossberg, "Transient Probe Spectra in Strongly Driven Atoms and their Dependence on Initial Atomic Conditions," Physical Review A33, 3956-3969 (1986) (additional support from NSF).
7. N. Lu, P.R. Berman, Y-S. Bai, J.E. Golub and T.W. Mossberg, "Time-dependent Spectrum of Resonance Fluorescence for Atoms Prepared in Pure Dressed States," Physical Review A34, 319-325 (1986) (additional support from NSF).
8. E.J. Robinson, "A General Solution to the Two-Level Problem," J. Phys. B18, 3687 (1985).
9. E.J. Robinson, "A general Solution to the Two-Level Problem: Further Results," J. Phys. B18, L657 (1985).
10. E.J. Robinson, "Adiabatic Switching and Long-Term Corrections to the Exponential Decay Law," Phys. Rev. A33, 1461 (1986).
11. E.J. Robinson, "Wigner's Infinite Product Representation of the R-Matrix: Rederivation, Interpretation and Potential Applications," Phys. Rev. A34, 1896 (1986).
12. E.J. Robinson "Quenched Exponential Decay". Phys. Rev. Letters 57, 1281 (1986).

BOOKS (AND SECTIONS THEREOF) SUBMITTED FOR PUBLICATION

BOOKS (AND SECTIONS THEREOF) PUBLISHED

1. P.R. Berman and R.G. Brewer "Modified Bloch Equations for Solids," in Laser Spectroscopy VII edited by T.W. Hänsch and Y.R. Shen (Springer-Verlag, Berlin, 1985) pp. 270-273. [This article also appears in Methods of Laser Spectroscopy edited by Y. Prior, A. Ben-Reuven, and M. Rosenbluh (Plenum, New York, 1986) pp. 205-208.]
2. F.P. Mattar, P.R. Berman, A.W. Matos, Y. Claude, C. Goutier and C.M. Bowden, "Three-level Superfluorescence" in Multiple-Photon Excitation and Dissociation of Polyatomic Molecules", edited by C.D. Cantrell (Springer-Verlag, Berlin, 1986), pp. 223-283 (additional support from ARO and NSF).

END

12-86

DTIC